

Horst Kächele (Ulm)

Phases of Research in Psychotherapy

abstract

Tel.: (0731) 502--5661 Fax: (0731)
e-mail: kaechele@sip.medizin.uni-ulm.de
<http://sip.medizin.uni-ulm.de/>

Psychotherapy research aims at the scientific evaluation of existing practice and at the discovery of new fields of application. The early phases of psychotherapy research were marked by scientific justification and societal legitimation. These questions changed with extension of possible indications, with growing differentiation of treatment procedures and with the progressive implementation of psychotherapy within the health system. The early approach "does psychotherapy work at all" has been replaced by the questions "how does psychotherapy work" and "what works for whom". Describing this phase model of psychotherapy research helps to better evaluate a variety of issues present in ongoing discussions.

Korrespondenz-Adresse:

Prof. Dr. med. Horst Kächele


Abteilung für Psychotherapie und Psychosomatische Medizin

Universität Ulm

Am Hochsraess 8, 89081 Ulm

Psychotherapy research at any given time has multiple functions and tasks to perform; it aims at the scientific evaluation of existing practice and at the discovery of new fields of application.

figure 1



The early period of psychotherapy research was marked by scientific justification and societal legitimation. Its questions changed with extension of possible indications, with growing differentiation of treatment procedures and with the progressive implementation of psychotherapy within the health system. The early approach "does psychotherapy work at all" has been replaced by the question "what works for whom" (Roth & Fonagy 1997) and "how does what kind of psychotherapy work".

Futhermore it has become obvious that the findings from systematic outcome research are directed at different audiences - e.g. at psychotherapists who conduct the treatment in question as well as to health professionals from related, often competitive disciplines. Research findings are addressed at those who benefit directly (e.g. patients or their relatives) as well as at those who fund the costs (e.g. insurance companies) or are responsible for adequate health policies (e.g. politicians, unions). The diverse groups may have totally different expectations (Strupp & Hadley 1977). Therefore outcome research has to provide a variety of information to satisfy the needs of the different interest groups.

The findings of many investigations consistently support the conclusion that overall psychotherapeutic treatments rightly have become an integral part of the medical system. This success opens a view for

raising new issues. We have to attend more care to the validity aspects of experimental process-outcome research demonstrating efficacy and of field studies on the treatment as applied in usual routine demonstrating effectiveness. It is on this distinction that this contribution hopes to make some contribution.

First I shall use a phase model of types of psychotherapy research that no necessary temporal order:

Phase I: descriptive study (*process research*)

Phase II: experimental analogue study (*process-outcome research*)

Phase III: clinical trial (*efficacy research*)

Phase IV: field study (*effectiveness research*)

Phase V: patient oriented treatment study (*efficiency research*)

Phase I descriptive study on the therapeutic process

We are now faced with the seeming paradox that, in spite of the overwhelming and certainly impressive evidence for the most frequently practiced forms of therapy, we are faced with many critical voices complaining that the many outcome studies have not contributed to a better understanding of therapeutic mechanisms (Grawe 1988). It is within this context that the very material of the therapeutic process is rediscovered and the detailed analysis of single cases once more achieves a prominent status (Dahl et al. 1988; Greenberg 1991; Greenberg & Pinsof 1986). This move entails increasingly focusing on microprocesses of the treatments requiring new assessment procedures and a better articulation of moment to moment events that may significantly influence treatment outcome (Kächele 1992).

Phase II experimental analogue study

The beginning of the experimental analogue derived from the classic studies of Greenspoon (1955) in the area of operant conditioning. Hans Strupp presumably was the first psychodynamic treatment researcher that used an experimental analogue approach in his studies on the therapist's technique (1960). The experimental mind of behaviorists have cared for a immense output on these kind of studies (Kiesler 1971); there have been really very few psychoanalytic minded researchers following the model set

by Strupp. The studies on the use of free association are a laudable exemption (see Heckmann et al. 1987).

Phase III: clinical trial (*efficacy research*)

This type of research has received utmost attention in the anglo-american worlds as thousands of so called Randomized Controlled Trial (RCT)'s have been performed. Much less have been performed in German speaking countries. Grawe et al. (1994) identified in 1985 more than 3.5 thousand of such studies. Using a number of sensible criteria they still ended up with 897 studies on adult real patients. The clinical trial consists in a comparison of two or more groups of patients in psychotherapy or in control conditions that are allotted by randomization to one of the conditions. Measurement of outcome is done using multiple perspectives and multiple measurements.

Ken Howard has recently given a succinct appreciation of the core features of RANDOMIZED CLINICAL TRIALS that emphasize the aspect of internal validity: He recommends:

Gather assessments of lots of plausible potential confounds (setting, treatment, therapist, and patient variables).

Stay clearly focused on the research question (in clinical research this question should be patient focused -- what is the best treatment for this (type of) patient?).

Probably the most well know study of this kind is the NIMH Treatment of Depression Collaborative Research Program (TDCRP) that has been described in great detail by Elkin 1994 in the fourth edition of the 'Handbook of Psychotherapy and Behavior Change' (Berhin & Garfield 1994) that set out to evaluate the value of two psychotherapy conditions (Cognitive-Behavioral Treatment (CBT) and Interpersonal Therapy (IPT) for the treatment of depression compared to a standard treatment with imipramine. Meanwhile this study has aroused many controversial discussions in the field on the validity of the findings (see Kächele 1996). I shall take up some remarks by Jacobson, a well know psychotherapy researcher, who has sharpened the debate regarding the usefulness of RCT for making decisions on clinical practice (Jacobsen 1995 in Kächele 1996):

First, it seems clear to me that clinical trials have generally provided optimal tests of the efficacy of CBT and other psychotherapies, and undoubtedly overestimate the value of these treatments as practiced by your typical clinician who is not rigorously trained, monitored, or supervised during the course of a trial.

Our research and that of others raises questions about the exportability of CBT as a treatment for depression into naturalistic settings, since competence seems to drift downward even among highly experienced therapists who were well-trained to a certain level of competence, unless supervision/calibration remains quite intensive throughout the trial. Any treatment which requires such intensive supervision, even with highly experienced and well-trained therapists, may not generalize well to typical practitioner settings where there is little training and certainly no supervision.

Second, the results in clinical trials are further skewed by subject selection procedures designed to homogenize the sample and detract from its representativeness to clinical practice. As Ellen Frank and Tracie Shea pointed out at the meeting, this subject selection practice cuts both ways. On the one hand, the single most often used basis for exclusion in depression trials is that subjects are not sufficiently depressed: so subjects treated in clinical trials are in all likelihood MORE depressed than those typically seeking therapy from practitioners. On the other hand, the exclusion of bipolar I and bipolar II patients, patients with dual diagnoses, including current episodes of substance abuse, panic disorder, eating disorders, and the like limit generalizability. But these two opposing processes (selecting for real major depressive disorder while at the same time screening out diagnostic complexity) result in a sample of patients that in some ways are easier to treat, and in other ways are harder to treat.

Third, even taking into account the highly select sample of patients selected for these trials, the efficacy of CBT, as well as other available pharmacological and psychosocial treatments, are exceedingly modest from the standpoint of clinical significance. Whether psychosocial treatments, including CBT, exceed rigorously constructed placebos is still very much open to question.

These comments were reiterated and reinforced by another prominent member of the US-scientific community, by Martin Seligman. Reporting on the findings of a consumer satisfaction study - the Consumer Reports - he concluded:

The efficacy study is the wrong method for empirically validating psychotherapy as it is actually done, because it omits too many crucial elements of what is done in the field (Seligman 1995, S.966).

To add to these critics Kächele et al. (1999) have compared the amount of treatment provided within such RCT by analyzing the studies reported on by Grawe et al (1994):

Cognitive-Behavioral Therapies

429 studies mean 11,2 sessions

434 studies mean 7, 9 weeks

Humanistic Therapies

70 studies mean 16,1 sessions

76 studies mean 11, 6 weeks

Psychodynamic Therapies

82 studies mean 27,6 sessions

80 studies mean 30,7 weeks

These findings again raise the question of the external validity of the RCT studies assuming that psychotherapy as practised in routine conditions deviates grossly from these findings.

Phase IV field study (*effectiveness research*)

Checking for facts describing what is routine practice in providing treatment in terms of time (number of sessions or month in treatment) Kächele et al. (1997) could identify three samples.

Sample A: 1800 psychodynamic oriented treatments delivered at the out-patient service of the department of psychotherapy at the Ulm University.

Sample B: 302 client-centered psychotherapies across Germany

Sample C: 496 behavioral therapies from the outpatient service of the department of clinical psychology at the Ruhr University Bochum

	mean	median	variation	
Sample A		22	6 - 1200	
Sample B	69,2, s 39,1	61	8 - 275	
Sample C	30,7, s 13,12	24	4 - 95	

The range of these treatments is considerable; it is most among psychoanalytic therapies. The graphical representations of these findings illustrate that the length of time in treatment replicates the findings of the dosage-effect relationship of Howard et al. (1987). Many treatments are terminated rather soon, a smaller number takes considerable time.

Investigations of these phenomena have far-reaching clinical implications because they correct the clinician's illusion (Vessey et al 1993) that he or she is treating an representative sample of patients. Epidemiological studies of the incidence and prevalence of psychosomatic or neurotic illnesses, of bodily dysfunctions or emotional disturbances, give an estimate of the need for services; the investigation of therapeutic practices yields an estimate of available resources to meet those needs; and, studies of the patterns of service utilization identify the constituencies served by the delivery system (Howard et al.1992).

Detailed field research describing what happens in treatment in many naturalistic dimensions on multiple cases combined with sophisticated outcome measurement supports finding specific treatments for patient groups with particular disorders. For many years substantial naturalistic research on specific groups of patients remained rare as the institutions had to take care of a broad spectrum of patients, which only sometimes allowed for the formation of groups homogeneous with respect to one disease. It took some time in psychotherapy to realize that this situation was not principally different from somatic medicine, where the development of multi-center studies led to progress because the desired homogeneity allowed better conclusions.

The multi-center study on the psychodynamic treatment of eating disorders that has been initiated by the Center for Psychotherapy Research in Stuttgart in 1992 includes a wide range of inpatient modalities all over Germany; it also has established the logistics of implementing the study in a considerable number of European countries. The German multi-center study is heavily supported by non-university institutions and offers also clinical exchange programs; so we feel that this type of research commitment may well turn out to be a prototype for the new look in psychotherapy research (Kächele et al. 1992). The results on a large sample of 1200 patients with anorexia and bulimia that are beginning to appear demonstrate that despite huge diversity of treatment modalities the

estimate of treatment success is not dependent on extent and duration of in-patient treatment (Kächele & project TR-EAT 1999; 2001).

Phase V patient focused treatment research

As the question of differential indication is what interests the clinician most (Kächele & Kordy 1997) this approach still does not answer the crucial question: does this particular treatment work for this patient ?


The main limitation of clinical trials and field studies is that they can only confirm the efficacy and effectiveness of treatments that have already been developed. They are not generative. Also, they do not speak to individual clients, as a more qualitative or single subject design would. At this point in the discussion of clinical research we have to ask: how does one study the potential impact of treatment on an individual client? We now know that predictive validity of clinicians' judgment on later outcome is not very impressive. Even if providing a standard interviewing procedure for predicting outcome therapists are performing very poor.

New approaches are in the process of being developed. First among these has been the work of Howard et al. (1996; 1997). It is obvious that this kind of research might turn out to be the really useful for the practicing therapist. Computer assisted feedback driven active quality management for psychotherapy provision most likely will turn out as a major service soon provided even via internet as the Stuttgart-Heidelberg Model (Kordy et al. 2001). is already developing.

References

- Dahl, H., Kächele, H., & Thomä, H. (Eds) *Psychoanalytic Process Research Strategies*. Berlin Heidelberg New York London Paris Tokyo: Springer.1988
- Grawe, K.: Zurück zur psychotherapeutischen Einzelfallforschung. *Z Klin Psychol*, 17 (1988b) 4
- Greenberg, L.: Research on the process of change. *Psychotherapy Research* 1 (1991) 3
- Greenberg L, Pinsof W (Eds) *The Psychotherapeutic Process: A Research Handbook*. Guilford Press, New York 1986.
- Greenspoon J (1955) The reinforcing aspect of two spoken sounds on the frequency of two responses. *American Journal of Psychology* 68:409-416
- Heckmann H, Hölzer M, Kächele H, Robben H (1987) Resistance and transference as two main constituents in an "experimental analogue" of free association. In: Huber W (Hrsg) *Progress in Psychotherapy Research*. Presses Universitaire de Louvain, Louvain-la -Neuve, S 582-593
- Howard KI., Kopta, S. M., Krause, M. S., & Orlinsky, D. E.: The dose-effect relationship in psychotherapy. *American Psychologist*, 41 (1986) 159
- Howard KI., Lueger RJ, Schank D.: The psychotherapy service delivery system. *Psychotherapy Research*, 2 (1992)
- Howard K, Martinovich Z (1997) An empirical basis for case assignment. In: Kächele H, Mergenthaler E, Krause R (Hrsg) *Psychoanalytic Priocess Research Strategies II: Twelve years later*. Ulmer Textbank: <http://sip.medizin.uni-ulm.de/psychotherapie>
- Howard K, Moras K, Brill P, Martinovich Z, Lutz W (1996) The evaluation of psychotherapy: Efficacy, effectiveness, patient progress. *Am Psychologist*
- Kächele H (1992) Narration and observation in psychotherapy research. Reporting on a 20 year long journey from qualitative case reports to quantitative studies on the psychoanalytic process. *Psychotherapy Research* 2:1-15
- Kächele H (Hrsg) (1996) *Letters from the SSCPnet*. Ulmer Textbank, Ulm
- Kächele, H. u. Mitarbeiter (1992). Planungsforum "Psychodynamische Therapie von Eßstörungen". *PPmP DiskJournal*. 42(3:1)
- Kächele H für die Studiengruppe MZ-ESS (1999) Eine multizentrische Studie zu Aufwand und Erfolg bei psychodynamischer Therapie von Eßstörungen: Studiendesign und erste Ergebnisse. *PPmP Psychother Psychosom med Psychol* 49: 100-108

- Kächele H, Kordy H, Richards M (2001) Therapy amount and outcome of inpatient psychodynamic treatment of eating disorders in Germany: Data from a mulitcenter study. *Psychotherapy Research* 11: 239-257
- Kächele H, Schulte W, Hillecke W Eckert J (2001) Zeit für Psychotherapie. in Vorbereitung
- Kächele, H., & Kordy, H. (1997). Outcome Research. In T. v. Uexküll (Ed.), *Psychosomatic Medicine* (pp. 290-299). New York: Urban & Schwarzenberg.
- Kiesler DJ (1971) Experimental designs in psychotherapy research. In: Bergin AE, Garfield SL (Hrsg) *Handbook of Psychotherapy and Behavior Change*. Wiley & Sons, 36-74
- Kordy H, Hannover W, Richard M (2001) Computer-assisted feedback-driven quality managment for psychotherapy. *J Cons Clin Psychol* 69: 173-183
- Newmann FL, Howard KI: Therapeutic effort, treatment outcome, and national health policy. *Amer Psychol* 41 (1986) 181
- Roth A, Fonagy P (1996) what works for whom ? - A critical review of psychotherapy research. Guilford Press, New York - London
- Sandell R, Blomberg J, Lazar A (1997) When reality doesn't fit the blueprint: doing research on psychoanalysis and long-term psychotherapy in a public health service program. *Psychother Research* 7: 333-344
- Seligman M (1994) What you can change and what you can't. The complete guide to successful self-improvement. A Knopf, New York
- Seligman MEP (1995) The effectiveness of psychotherapy. The *Consumer Reports* Study. *Am Psychologist* 50:965-974
- Strupp HH (1960) *Psychotherapists in Action: Explorations of the Therapist's Contribution to the Treatment Process*. Grune & Stratton, New York
- Strupp HH, Hadley SW: A tripartite model of mental health and therapeutic outcome. *Amer. Psychol.* 32 (1977) 187
- Vessey JT, Howard KI, Lueger RJ, Kächele H, Mergenthaler E (1994) The Clinician's Illusion and the Psychotherapy Practice: An Application of Stochastic Modeling. *J Couns Clin Psychol* 62: 679-685



Phases of Research

Phase I: descriptive study

(process research)

Phase II: experimental analogue study

(process-outcome research)

Phase III: clinical trial

(efficacy research)

Phase IV: field study

(effectiveness research)

Phase V: patient oriented treatment study

(efficiency research)

The efficacy study is the wrong method for empirically validating psychotherapy as it is actually done, because it omits too many crucial elements of what is done in the field

(Seligman 1995, S.966).